

Effect of three drying methods on volatile oil of turmeric

ABSTRACT: **OBJECTIVE** To analyze the effects of sun drying, blast drying and [microwave drying equipment](#) on the volatile oil of *Curcuma longa* Wenyujin *Rhizoma Concisum*. **METHODS** The volatile oil was extracted by steam distillation after *Curcuma* *Curcuma* was dried, analyzed and identified by GC-MS, and then the relative contents of each component were determined.

Results Twenty-six components were identified in volatile oil of sun dried products, 29 in blast dried products and 28 in microwave dried products. The relative content of zedoary Dione was the highest among the three dried products, especially in sun dried products, but it did not contain neozedoary dione.

CONCLUSION After drying by three methods, the components of volatile oil from *Curcuma longa* L. tablets are basically the same, but the relative contents are different. Under the conditions of blast drying and microwave drying, the complex isomerization of curcumadione and neocurcumadione may occur, which changes the relative content of curcumadione.

Key words: [microwave drying of traditional Chinese medicine](#); turmeric; volatile oil; sun drying; blast drying; microwave drying; steam distillation; GC-MS



Curcuma officinalis is a dry rhizome of *Curcuma wenyujin*, which was first recorded in the Compendium of Materia Medica. It has warm nature, bitter taste, and belongs to the spleen and liver meridians. It has the function of breaking blood, promoting qi, regulating menstruation and relieving pain. It mainly treats thoracic and hypochondriac stabbing pain, chest stuffiness and heartache, dysmenorrhea, drop and paralysis, etc.

It is one of the freshly cut pieces of Chinese herbal medicine. It is digged, washed, and removed fibrous roots after the stem and leaf withered in winter. It is usually dried by sun or blast. In recent years, microwave drying has also been applied to the processing of its origin. The herb mainly contains volatile components, but there are no reports about the effects of different drying methods on the volatile components of turmeric tablets.

Therefore, this experiment used GC-MS method to analyze the changes of volatile oil composition after fresh-cut turmeric tablets were processed by blast drying, microwave drying

and sun drying, and the effects of different drying methods on them were analyzed.

The volatile oil of *Curcuma longa* is mainly composed of sesquiterpenes and a few monoterpenes. It has many pharmacological activities, such as anti-tumor, anti-inflammatory, anti-oxidation, reducing blood lipid, improving blood rheology, analgesia and so on. The former is the basis of its main active substance.

At present, the related pharmacological studies mainly focus on sesquiterpenes, such as the anti-tumor, anti-bacterial, anti-virus, antitussive and antiasthmatic effects of gemcitrone, the anti-cancer and anti-virus effects of neozedoary ketone, and the anti-tumor, anti-coagulation, anti-thrombosis, anti-inflammatory, analgesic and liver protection effects of zedoary ketone.

It was found that the main components of volatile oil under sun drying, blast drying and microwave drying were mainly anti-tumor, anti-inflammatory and anti-virus. That is to say, the pharmacological effects of the three components may not change significantly, and bioequivalence evaluation is needed.

In summary, the composition and relative content of volatile oil of *Curcuma longa* L. under the three drying methods are basically the same, but there are some differences in trace components. Therefore, through the comparative analysis of the three, this experiment can provide theoretical basis for the study of processing and drying methods and standardized production of turmeric slices.

